

# Hydroxychloroquine and covid-19

Neeraj Sinha ,<sup>1</sup> Galit Balayla<sup>2</sup>

<sup>1</sup>Pulmonary Transplant, University of Miami Miller School of Medicine, Miami, Florida, USA

<sup>2</sup>General Medicine, Central University of Venezuela, Caracas, Venezuela

## Correspondence to

Dr Neeraj Sinha, Pulmonary Transplant, University of Miami Miller School of Medicine, Miami, FL 33136-1015, USA; nsinhamd@yahoo.com

Received 30 March 2020  
Accepted 1 April 2020

## ABSTRACT

Hydroxychloroquine and chloroquine are medications that have been used for a long time. Their most common use is for the treatment and prophylaxis of malaria. However, these antimalarial drugs are known to also have anti-inflammatory and antiviral effects and are used for several chronic diseases such as systemic lupus erythematosus with low adverse effects. The antiviral action of hydroxychloroquine and chloroquine has been a point of interest to different researchers due to its mechanism of action. Several in vitro studies have proven their effectiveness on severe acute respiratory syndrome virus and currently both in vitro and in vivo studies have been conducted on 2019 novel coronavirus (covid-19). The purpose of this article is to review the history and mechanism of actions of these drugs and the potential use they can have on the current covid-19 pandemic.

## INTRODUCTION

Antimalarial drugs such as chloroquine and hydroxychloroquine have been in the market for over a century. These drugs have been used not only for malaria but also for several rheumatic diseases given their anti-inflammatory properties, affordability and the fact that they have shown a good safety profile. Knowing that these drugs have proven effective for a wide range of diseases has made several researchers wonder about their use in other areas such as cancer and viral infections. This is why, in the midst of a global pandemic, the question of antimalarial use in treatment and prophylaxis of covid-19 has been raised.

## HISTORICAL USES AND SIDE EFFECTS

Uses of Cinchona bark for a 'febrile disease' have been reported since 1630 in Lima, Peru. In 1894, Dr J.F. Payne was the first one to administer it at St. Thomas Hospital in London for something other than malaria. He used it in the treatment of a rash that we now can infer was a lupus rash.<sup>1,2</sup> Chloroquine was discovered in 1939 by the Germans. It substituted quinacrine by the end of World War II for the treatment of malaria proving to be safer and more effective.<sup>3</sup> During the war, soldiers were given prophylactic antimalarial drugs which incidentally improved lupus erythematosus rashes and arthritis.<sup>4</sup> Hydroxychloroquine was introduced to the market in 1955, differing from chloroquine by a hydroxyl group that makes it safer. These drugs have also proven to be effective in other inflammatory diseases such as rheumatoid arthritis,<sup>4,5</sup> and in 1989, Geser *et al* described the decreased incidence of Burkitt lymphoma in patients treated for malaria prophylactically with chloroquine.<sup>6</sup> These different uses are summarised by Ben Zvi *et al*.<sup>7</sup>

Hydroxychloroquine and chloroquine are relatively safe drugs. Their most common side effects include gastrointestinal symptoms, pruritus and dermatological changes that can occur in up to 10% of the patients.<sup>8</sup> The most severe side effects have low incidence. They include neuromyopathy of proximal muscles, cardiotoxicity and irreversible retinopathy. The latter is well documented in long-term users with high doses. This can be prevented or controlled with dose calculation based on body weight, reducing the dose after 5 years of use and routine ophthalmologic evaluation.<sup>9</sup> On the other hand, the neuromuscular changes can slowly improve by discontinuing the medication. The cardiotoxicity may include QT prolongation syndrome, especially in patients with renal or hepatic dysfunction. Overdose toxicity has also been observed between 1 and 3 hours after a high-dose intake, especially with chloroquine. Toxic effects have occurred when dosing at 20 mg/kg and have been fatal at over 30 mg/kg. The overdose presents with visual changes, nausea, hypokalaemia, drowsiness, shock, convulsions and even death. Diazepam has been used as an antidote. Hydroxychloroquine overdose is rare and the dose has not been well established.<sup>10</sup>

## ANTI-INFLAMMATORY EFFECT

The effect of hydroxychloroquine and chloroquine on the immune system has been well established. Several of their anti-inflammatory mechanisms have been recognised, for example, interference with lysosomal acidification and antigen presentation,<sup>11</sup> inhibition of phospholipase A2,<sup>12</sup> absorption and blocking UV light cutaneous reactions,<sup>13</sup> binding and stabilising DNA,<sup>14</sup> inhibition of toll-like receptor signals, inhibition of T and B cell receptors, and especially, decreasing cytokine production by macrophages such as interleukin (IL)-1 and IL-6.<sup>7,11</sup> Interaction with toll-like receptors and T cell receptors makes hydroxychloroquine effective in different sites of the signalling cascade in the inflammatory response. These interactions prevent autoimmunity without immunosuppressing the patient.<sup>15</sup> Sperber *et al* also demonstrated the inhibition of IL-1 and IL-6 in T cells and monocytes<sup>11</sup> and several other studies have demonstrated tumour necrosis factor (TNF)-alpha inhibition or attenuated inflammatory effect by hydroxychloroquine.<sup>16-18</sup> The effective inhibition of inflammatory cytokines such as IL-6, IL-1 and TNF-alpha decreases tissue damage and endothelial inflammation thus preventing initiation and propagation of autoimmune inflammation.<sup>19</sup> This cytokine inhibition is of great importance at this time, since it has been demonstrated that several viruses upregulate the expression of IL-1, IL-6 and TNF-alpha



© Author(s) (or their employer(s)) 2020. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Sinha N, Balayla G. *Postgrad Med J* Epub ahead of print: [please include Day Month Year]. doi:10.1136/postgradmedj-2020-137785

**Table 1** Ongoing clinical trials for Hydroxychloroquine on COVID-19.

Name	Identifier	Location	1. Type of study 2. What to measure	Treatment arm	Active comparator	Placebo arm
Randomized Controlled Clinical Trials of Lopinavir/Ritonavir or Hydroxychloroquine in Patients With Mild Coronavirus Disease (COVID-19)	NCT04307693	Sung-Han Kim, Asan Medical Center, Seoul, South Korea	1. Multicentre, open-labelled, randomised clinical trial 2. Viral load	Lopinavir/ritonavir 200 mg/100 mg two tablets by mouth, every 12 hours for 7–10 days	Hydroxychloroquine 200 mg two tablets by mouth, every 12 hours for 7–10 days	No lopinavir/ritonavir and hydroxychloroquine
Hydroxychloroquine Post Exposure Prophylaxis for Coronavirus Disease (COVID-19)	NCT04318444	Elizabeth Oelsner, Columbia University, NY, USA	Parallel assignment (randomised)	Hydroxychloroquine two tablets (400 mg) twice daily on day 1; for days 2–5, they will be instructed to take one tablet (200 mg) twice daily		Two tablets (400 mg) twice daily on day 1; for days 2–5, they will be instructed to take one tablet (200 mg) twice daily
Hydroxychloroquine Chemoprophylaxis in Healthcare Personnel in Contact With COVID-19 Patients (PHYDRA Trial)	NCT04318015	National Institute of Respiratory Diseases, Mexico City, Mexico	1. Triple blinded, randomised controlled trial 2. Symptomatic covid-19 infection rate	1. High risk: hydroxychloroquine 200 mg per day for 60 days 2. Low risk: hydroxychloroquine 200 mg per day for 60 days		1. High risk: placebo tablet per day for 60 days 2. Low risk: placebo tablet per day for 60 days
Chloroquine Prevention of Coronavirus Disease (COVID-19) in the Healthcare Setting (COPCOV)	NCT04303507	Oxford University, Oxford, UK	1. Parallel assignment 2. Number of symptomatic covid-19 infections	Chloroquine A loading dose of 10 mg base/kg followed by 150 mg daily (250 mg chloroquine phosphate salt) will be taken for 3 months Hydroxychloroquine (400 mg 2 times per day, 12/12 hours)+azithromycin (500 mg once a day)		Placebo
Safety and Efficacy of Hydroxychloroquine Associated With Azithromycin in SARS CoV-2 Virus (Alliance COVID-19 Brasil II)	NCT04321278	Hospital Israelita Albert Einstein, Sao Paulo, Brazil	1. Parallel assignment 2. Evaluation of clinical status		Hydroxychloroquine (400 mg 2 times per day, 12/12 hours)	
Post-exposure Prophylaxis for SARS-Coronavirus-2	NCT04308668	University of Minnesota, Minnesota, USA	1. Parallel assignment 2. Incidence of disease	Hydroxychloroquine 200 mg tablet; 800 mg orally once, followed in 6–8 hours by 600 mg, then 600 mg once a day for four consecutive days		Placebo four placebo tablets once, followed in 6–8 hours by three tablets, then three tablets once a day for four consecutive days

in vitro, suggesting the effectiveness of hydroxychloroquine in viral infections.

### INFLAMMATORY RESPONSE IN VIRAL INFECTIONS

The role of inflammatory cytokines in viral disease is not well understood; nevertheless, numerous hypotheses have been proposed to describe the viral inflammatory response. It is believed that IL-6 can have both inflammatory and anti-inflammatory effect in the body depending on which signal is activated: IL-6 trans signalling is inflammatory while regular IL-6 signalling is anti-inflammatory.<sup>20</sup> However, several studies point to negative implications of IL-6 involvement in the immune response to viral infections. Three mechanisms have been suggested: (1) in vitro secretion of IL-6 by toll-like receptor activation causes an inhibition of CD8 T cell response, which is confirmed by an in vivo blockage of IL-6 with monoclonal antibodies. This blockage causes reduction in viral loads and increases production of interferon (IFN)-gamma<sup>17</sup>; (2) the synergistic interaction of IL-6 and IL-17 generates persistence of viral infection and worsens clinical outcomes; and (3) it is suggested that IL-6 upregulates PD-1 and PDL-1. Normally PD-1 and PDL-1 interaction prevents autoimmunity through programmed cell death. However, during a viral infection, it alters the immune response against the virus reducing CD8+ cytolytic function that is worsened by the upregulation from IL-6.<sup>21</sup> Many studies have confirmed the increased levels of IL-6 and TNF-alpha in viral infections such as hepatitis C virus, HIV, hepatitis B virus, influenza virus, chikungunya virus and severe acute respiratory syndrome virus (SARS-CoV).<sup>20–24</sup>

### HYDROXYCHLOROQUINE/CHLOROQUINE ANTIVIRAL EFFECT

The effect of hydroxychloroquine and chloroquine on viral replication goes beyond cytokine inhibition. These medications are weak bases that can affect acid vesicles and inhibit several enzymes. This characteristic allows them to inhibit the viral entry to the cell when the endocytosis is pH dependent. It also inhibits glycosyl-transferases, viral post-translational modifications and replication of some viral families. The antiretroviral effect has been considered to be caused by the inhibition of viral glycosylation, a major antiviral mechanism of these drugs.<sup>24</sup> It was also recently described that chloroquine might inhibit quinone reductase-2, an enzyme involved in sialic acid biosynthesis. If this were to be true, it would explain further its effect on HIV, SARS and *orthomyxoviruses* because sialic acid is present on HIV-1 glycoproteins, SARS angiotensin-converting enzyme 2 (ACE2 receptor and *orthomyxovirus* receptors).<sup>25 26</sup>

Severe acute respiratory syndrome (SARS), better known as SARS-CoV, was a novel *coronavirus* that emerged in China in November 2002. It rapidly spread to more than 30 countries, causing 8096 cases and 774 deaths. The syndrome is described with fever, chills, malaise, cough and dyspnoea that can progress to respiratory failure.<sup>27</sup> Savarino *et al* were the first ones to hypothesise that hydroxychloroquine and chloroquine might be helpful in treatment of SARS, since endocytosis might be involved in viral entry to the cell and there was an important immune response that was causing clinical worsening, probably due to inflammatory cytokines such as TNF-alpha and IL-6.<sup>24</sup> This was later confirmed by two different in vitro studies. Kayaerts *et al* demonstrated the inhibition of SARS-CoV by chloroquine in Vero E6 cells at different postinfection times.<sup>28</sup> Vincent *et al* proved the effective dose-dependent inhibition of the virus in Vero E6 cells immediately after viral absorption and also 3–5 hours after absorption. They also showed

that pretreated cells with chloroquine were refractory to the virus and revealed an impairment of terminal glycosylation of ACE2 receptor, decreasing viral-receptor affinity and therefore reducing the initiation of the infection. This experiment can illustrate the possibility of using hydroxychloroquine for coronavirus prophylaxis the same way as in malaria.<sup>29</sup>

### THE CURRENT CORONAVIRUS OUTBREAK

In December 2019, a coronavirus outbreak started in Wuhan, China. The viral infection presented with high fever, dry cough, headache, dyspnoea and diarrhoea. The first cases that showed these symptoms were all associated with a food market in the city of Wuhan. Since then, there have been several cases that progressed to respiratory failure, with an estimated mortality rate of 2%. According to the WHO, to this date, 23 March 2020, there are 332 935 confirmed cases, 14 510 deaths and 190 affected countries.<sup>30</sup> In order to understand this outbreak, the genetic sequencing of the virus has been extracted and is now known to share 79% of its genome with SARS-CoV from the Coronaviridae family. However, it is sufficiently different from it to be considered a *novel* virus. The WHO renamed it **covid-19** (SARS-CoV-2) in February 2020 and declared it a **Pandemic**. It has also been demonstrated that this novel virus not only shares sequencing with SARS-CoV but also uses the same cell entry ACE2 receptor.<sup>31 32</sup> Even though the *novel* virus shares genomic sequencing with SARS-CoV, data show that their shedding patterns in infected patients differ substantially, making covid-19's more similar to the pattern of influenza virus. Furthermore, the viral load of covid-19 has been similar in asymptomatic patients and symptomatic patients, thus, making the virus transmissible in early stages of the infection. On the contrary, transmission of SARS-CoV occurred within days of initial symptoms, making the epidemiological strategies different for the two infections.<sup>33</sup>

### USES OF HYDROXYCHLOROQUINE AND CHLOROQUINE IN COVID-19 (SARS-COV-2)

The worldwide effects of the SARS-CoV-2 pandemic has been unparalleled and it has prompted the scientific community to consider all possible solutions. Due to covid-19 similarities to SARS-CoV, several researchers have proposed the use of hydroxychloroquine and chloroquine on the novel virus.<sup>34–36</sup> Wang *et al* tested the effect of several Food and Drug Administration-approved antiviral drugs on the virus in vitro. Remdesivir showed post entry blockage of viral infection with an effective concentration at 50% of  $EC_{50}=0.77\ \mu\text{M}$  and a cytotoxic concentration of 50% of  $CC_{50}>100\ \mu\text{M}$ . Chloroquine was found to have an effective concentration at 50% of  $EC_{50}=1.13\ \mu\text{M}$ , a cytotoxic concentration at 50% of  $CC_{50}>100\ \mu\text{M}$  and an effective concentration at 90%  $EC_{90}$  of  $6.90\ \mu\text{M}$ . Chloroquine showed effectiveness at an entry and post entry level, while remdesivir was only effective at a post entry level. This further suggests the possible prophylactic use of chloroquine on SARS-CoV-2.<sup>37</sup> Yao *et al* also tested the effect of hydroxychloroquine and chloroquine in vitro. They divided the experiment into two phases: treatment study and prophylaxis study. In the treatment study, the  $EC_{50}$  values for chloroquine were 23.90 and  $5.47\ \mu\text{M}$  at 24 and 48 hours, respectively, and the  $EC_{50}$  values for hydroxychloroquine were 6.14 and  $0.72\ \mu\text{M}$  at 24 and 48 hours, respectively. In the prophylaxis study, the  $EC_{50}$  values for chloroquine were  $>100$  and  $18.01\ \mu\text{M}$  at 24 and 48 hours, respectively, and the  $EC_{50}$  values for hydroxychloroquine were 6.25 and  $5.85\ \mu\text{M}$  at 24 and 48 hours, respectively. They concluded that hydroxychloroquine

is more effective in vitro than chloroquine for both prophylaxis and treatment.<sup>38</sup>

The promising results of the in vitro experiments have led to the creation of multiple clinical trials to further investigate the chloroquine effect on covid-19. Currently, there are two trials with available data. Gao *et al* demonstrated the superiority of chloroquine over the control treatment in more than 100 patients regarding inhibition of pneumonia exacerbation, improvement in lung imaging findings, promoting a virus negative conversion and shortening the disease course in more than 10 hospitals in China.<sup>39</sup> Gautret *et al* treated 20 patients with hydroxychloroquine and compared the results with 16 controls in France. They used PCR to measure the viral load on day 3, 4, 5 and 6 of postinclusion. The treatment group had a higher age mean, but no difference in gender was made between the two groups. Asymptomatic patients and patients with both lower and upper respiratory tract infections were treated. They concluded that hydroxychloroquine was effective in viral load reduction. The results on day 3 indicated that 50% of the hydroxychloroquine-treated patients had a viral load reduction with a  $p=0.005$ ; on day 4, it showed a 60% reduction with a  $p=0.04$ ; on day 5, a 65% reduction with a  $p=0.006$ ; and on day 6, 70% of the patients showed viral load reduction with a  $p=0.001$ . Furthermore, they described the synergistic effect of azithromycin when using it alongside hydroxychloroquine in decreasing the viral load. The dual treatment showed 100% decrease on the viral load with a  $p<0.001$  by day 6, while hydroxychloroquine alone showed a 70% decrease.<sup>40</sup>

### ONGOING STUDIES AND GUIDELINES

Several randomised clinical trials regarding antimalarial drug use in covid-19 have been proposed. On the US National Institute of Health's (NIH) medical library, there are already 143 trials on covid-19 registered from all over the world and 14 of them are specifically about the use of antimalarial drugs on the virus. These trials aim to demonstrate the effectiveness of hydroxychloroquine and chloroquine in pre-exposure and postexposure prophylaxis and treatment. Table 1 enlists six of these studies, but further information can be found at the NIH web page.<sup>41</sup> Moreover, on the Chinese clinical trial entry there are over 500 studies enrolled and 16 of them are about hydroxychloroquine and chloroquine use on covid-19. This information can be found on their web page.<sup>42</sup> Multiple institutions worldwide are already using hydroxychloroquine in their treatment guidelines, including the Centers for Disease Control and Prevention (CDC),<sup>43</sup> pointing to the relevance of this drug in the current pandemic.

### CONCLUSION

In the midst of a pandemic with a high mortality rate, the collapsing of health systems and the devastating effects on the economy worldwide, a fast solution is crucial. Preventive medicine is key in any pandemic. Preventing cases rather than treating them would decrease world morbimortality. Vaccinations have been the centre of prophylactic measures on infectious diseases. However, because of the novelty of the covid-19 virus, the undertermined immune reaction and its rapid spread, the creation of a successful vaccine for this virus in the short term is uncertain. Current measures such as self-quarantine and social distancing are recommended by institutions like the WHO and the CDC of Atlanta.

Covid-19 pandemic presents the classic conflict between the clinical bedside medicine and academic medicine. While clinical

### Main messages

- ▶ Hydroxychloroquine has shown several antiviral mechanisms, including the inhibition of inflammatory cytokines such as IL-1, IL-6 and TNF-alpha.
- ▶ The effect of hydroxychloroquine on SARS-CoV-2 (covid-19) has been studied in vitro, demonstrating its pre-entry result, probably due to the inhibition of the virus ACE2 receptor and the viral inhibition post-entry. Also, in vivo studies have demonstrated clinical improvement and decrease in the viral load.
- ▶ Several double-blinded, randomised, placebo-controlled clinical trials are being conducted to further investigate the effects of hydroxychloroquine on covid-19.

### Current research questions

- ▶ Should hydroxychloroquine be used as primary prophylaxis on covid-19?
- ▶ Can hydroxychloroquine be used as post-exposure prophylaxis on covid-19?
- ▶ Is hydroxychloroquine better than placebo in the treatment of covid-19?
- ▶ Is hydroxychloroquine better than other antiviral drugs such as ritonavir or lopinavir in the treatment of covid-19?

### Key references

- ▶ Andrea Savarino, John Boelaert, Antonio Cassone, *et al*. Effects of chloroquine on viral infections: an old drug against today's diseases. *Lancet Infect Dis* 2003;3:P722–7. DOI: [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(03\)00806-5/fulltext#sec57059262e394](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(03)00806-5/fulltext#sec57059262e394).
- ▶ Andrea Savarino, Livia Di Trani, Isabella Donatelli, *et al*. New insights into the antiviral effects of chloroquine. 2006;6:P67–9. DOI: [https://www.thelancet.com/journals/laninf/article/PIIS1473-3099\(06\)70361-9/fulltext](https://www.thelancet.com/journals/laninf/article/PIIS1473-3099(06)70361-9/fulltext).
- ▶ Xueting Yao, Fei Ye, Miao Zhang, *et al*. In vitro antiviral activity and projection of optimised dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020;pii: ciaa237. doi: 10.1093/cid/ciaa237. <https://academic.oup.com/cid/advance-article/doi/10.1093/cid/ciaa237/5801998>.
- ▶ Jianjun Gao, \* Zhenxue Tian, Xu Yang. Breakthrough: Chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *BioScience Trends* 2020;14(1):72–3. DOI: 10.5582/bst.2020.01047. [https://www.jstage.jst.go.jp/article/bst/14/1/14\\_2020.01047/\\_pdf/-char/en](https://www.jstage.jst.go.jp/article/bst/14/1/14_2020.01047/_pdf/-char/en).
- ▶ Philippe Gautret, Jean-Christophe Lagier, Philippe Parola, *et al*. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an openlabel non-randomised clinical trial. 2020. <https://doi.org/10.1016/j.ijantimicag.2020.105949>. <https://www.sciencedirect.com/science/article/pii/S0924857920300996>.



## Self-assessment questions

- Which of the following adverse effects are caused by hydroxychloroquine?
  - Photosensitivity
  - QT prolongation
  - Retinopathy
  - Headache
- Which of the following mechanisms of action have been identified in hydroxychloroquine and chloroquine?
  - Inhibit DNA translation
  - Blocking UV cutaneous reactions
  - Inhibit inflammatory cytokines (IL-1/IL-6)
  - Block acetylcholine receptors
  - Block ACE2 receptors
  - Affect proteolysis
  - Affects glycosylation
- Which of the following statements are true about SARS-CoV?
  - It was a pandemic
  - Hydroxychloroquine was proven to be effective at a pre-entry and post-entry level
  - It started in the Middle East
- Which of the following statements are true about SARS-CoV-2 (covid-19)
  - It started at a food market in China
  - Has a mortality rate of 2%
  - It has not been declared a pandemic
  - It has the same cell receptor as SARS-CoV known as ACE2 receptor
  - The shedding pattern is more similar to SARS-CoV than influenza
- Which of the following statements are true about hydroxychloroquine effect on covid-19
  - It prevents the virus to enter the cell through ACE2 receptors
  - Has not shown a decreasing effect of the viral load
  - There are several countries currently using hydroxychloroquine to treat covid-19

bedside medicine incorporates evidence-based medicine, it also allows for prescription of the therapies which may not be based on the most rigorous clinical evidence, but based on biological plausibility, in vitro preclinical data or limited clinical evidence. In a setting like this, many hospitals have already adopted hydroxychloroquine-based therapies for covid-19 under the auspices of clinical protocols, the key ingredient of the clinical protocols being a mechanism to collect data prospectively so that we can learn as much as possible about the safety and efficacy of hydroxychloroquine therapy. In order to obtain the highest grade of evidence, double-blinded, randomised, placebo-controlled trials have also been started by a few other groups to determine the effectiveness of hydroxychloroquine on covid-19. The authors of this review article do not make any strong recommendations about which one of the above pathways to take. Well-established safety profile, encouraging but limited in vitro and in vivo data supporting benefit, low cost and easy availability, makes a clinical protocol pathway an understandable one for this novel disease with high mortality. We encourage the ongoing investigation on this area to have a clearer guideline to prevent and treat this disease.

**Twitter** Neeraj Sinha @md\_neeraj

**Contributors** Both authors contributed equally to the preparation of manuscript.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; internally peer reviewed.

## ORCID iD

Neeraj Sinha <http://orcid.org/0000-0002-6028-3271>

## REFERENCES

- Mates M, Nesher G, Zevin S. [Quinines--past and present]. *Harefuah* 2007;146:560–2.
- D'Cruz D. Lupus and the art of clinical medicine. *Rheumatology* 2015;54:i6–7.
- Tanenbaum L, Tuffanelli DL. Antimalarial agents. chloroquine, hydroxychloroquine, and quinacrine. *Arch Dermatol* 1980;116:587–91.
- Page F. Treatment of lupus erythematosus with mepacrine. *Lancet* 1951;2:755–8.
- Clark P, Casas E, Tugwell P, et al. Hydroxychloroquine compared with placebo in rheumatoid arthritis. A randomized controlled trial. *Ann Intern Med* 1993;119:1067–71.
- Geser A, Brubaker G, Draper CC. Effect of a malaria suppression program on the incidence of African Burkitt's lymphoma. *Am J Epidemiol* 1989;129:740–52.
- Ben-Zvi I, Kivity S, Langevitz P, et al. Hydroxychloroquine: from malaria to autoimmunity. *Clinic Rev Allerg Immunol* 2011;42.
- Wallace DJ. Antimalarial drugs in the treatment of rheumatic diseases, 2020. Available: [https://www.uptodate.com/contents/antimalarial-drugs-in-the-treatment-of-rheumatic-disease?search=hydroxychloroquine&source=search\\_result&selectedTitle=2~148&usage\\_type=default&display\\_rank=1#H7](https://www.uptodate.com/contents/antimalarial-drugs-in-the-treatment-of-rheumatic-disease?search=hydroxychloroquine&source=search_result&selectedTitle=2~148&usage_type=default&display_rank=1#H7)
- Abdulaziz N, Shah AR, McCune WJ. Hydroxychloroquine: balancing the need to maintain therapeutic levels with ocular safety: an update. *Curr Opin Rheumatol* 2018;30:249–55.
- Taylor WRJ, White NJ. Antimalarial drug toxicity: a review. *Drug Saf* 2004;27:25–61.
- Sperber K, Quraishi H, Kalb TH, et al. Selective regulation of cytokine secretion by hydroxychloroquine: inhibition of interleukin 1 alpha (IL-1-alpha) and IL-6 in human monocytes and T cells. *J Rheumatol* 1993;20:803–8.
- Löffler BM, Bohn E, Hesse B, et al. Effects of antimalarial drugs on phospholipase A and lysophospholipase activities in plasma membrane, mitochondrial, microsomal and cytosolic subcellular fractions of rat liver. *Biochim Biophys Acta* 1985;835:448–55.
- Lester RS, Burnham TK, Fine G, et al. Immunologic concepts of light reactions in lupus erythematosus and polymorphous light eruptions. I. The mechanism of action of hydroxychloroquine. *Arch Dermatol* 1967;96:1–10.
- Cohen SN, Yielding KL. Spectrophotometric studies of the interaction of chloroquine with deoxyribonucleic acid. *J Biol Chem* 1965;240:3123–31.
- Shukla AM, Wagle Shukla A, Shukla AW. Expanding horizons for clinical applications of chloroquine, hydroxychloroquine, and related structural analogues. *Drugs Context* 2019;8:2019-9-1.
- van den Borne BE, Dijkman BA, de Rooij HH, et al. Chloroquine and hydroxychloroquine equally affect tumor necrosis factor-alpha, interleukin 6, and interferon-gamma production by peripheral blood mononuclear cells. *J Rheumatol* 1997;24:55–60.
- Li R, Lin H, Ye Y, et al. Attenuation of antimalarial agent hydroxychloroquine on TNF- $\alpha$ -induced endothelial inflammation. *Int Immunopharmacol* 2018;63:261–9.
- Karim Sacre,<sup>1,2</sup> Lindsey A Criswell,<sup>3</sup> and Joseph M McCune. 2012. Hydroxychloroquine is associated with impaired interferon-alpha and tumor necrosis factor-alpha production by plasmacytoid dendritic cells in systemic lupus erythematosus. *Arthritis Res Ther* 2012;14:R155.
- Moudgil KD, Choubey D. Cytokines in autoimmunity: role in induction, regulation, and treatment. *J Interferon Cytokine Res* 2011;31:695–703.
- Wu W, Dietze KK, Gibbert K, et al. TLR ligand induced IL-6 counter-regulates the anti-viral CD8(+) T cell response during an acute retrovirus infection. *Sci Rep* 2015;5:10501.
- Velazquez-Salinas L, Verdugo-Rodriguez A, Rodriguez LL, et al. The role of interleukin 6 during viral infections. *Front Microbiol* 2019;10:1057.
- Breen EC, Rezaei AR, Nakajima K, et al. Infection with HIV is associated with elevated IL-6 levels and production. *J Immunol* 1990;144:480–4.
- Lan T, Chang L, Wu L, et al. IL-6 plays a crucial role in HBV infection. *J Clin Transl Hepatol* 2015;3:271–6.
- Savarino A, Boelaert JR, Cassone A, et al. Effects of chloroquine on viral infections: an old drug against today's diseases? *Lancet Infect Dis* 2003;3:722–7.
- Kwiek JJ, Haystead TAJ, Rudolph J. Kinetic mechanism of quinone oxidoreductase 2 and its inhibition by the antimalarial quinolones. *Biochemistry* 2004;43:4538–47.
- Savarino A, Di Trani L, Donatelli I, et al. New insights into the antiviral effects of chloroquine. *Lancet Infect Dis* 2006;6:67–9.
- Kenneth McIntosh MD. Severe acute respiratory syndrome (SARS), 2020. Available: <https://www.uptodate.com/contents/severe-acute-respiratory-syndrome-sars?search=>

- sars&source=search\_result&selectedTitle=1~29&usage\_type=default&display\_rank=1#H12
- 28 Keyaerts E, Vijgen L, Maes P, *et al.* In vitro inhibition of severe acute respiratory syndrome coronavirus by chloroquine. *Biochem Biophys Res Commun* 2004;323:264–8.
  - 29 Vincent MJ, Bergeron E, Benjannet S, *et al.* Chloroquine is a potent inhibitor of SARS coronavirus infection and spread. *Virology* 2005;2:69.
  - 30 World Health Organization. Novel coronavirus (2019-nCoV) technical guidance. Available: <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/technical-guidance>; <https://www.who.int/emergencies/diseases/novel-coronavirus-2019>
  - 31 Lu R, Zhao X, Li J, *et al.* Genomic characterisation and epidemiology of 2019 novel coronavirus: implications for virus origins and receptor binding. *Lancet* 2020;395:565–74.
  - 32 Zhou P, Yang X-L, Wang X-G, *et al.* A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* 2020;579:270–3.
  - 33 Zou L, Ruan F, Huang M, Ruan, and Dr. Huang. 2020. SARS-CoV-2 viral load in upper respiratory specimens of infected patients. *N Engl J Med* 2020;382:1177–9.
  - 34 Padmanabhan S. Potential dual therapeutic approach against SARS-CoV-2/COVID-19 with nitazoxanide and hydroxychloroquine 2020.
  - 35 Touret F, de Lamballerie X. Of chloroquine and COVID-19. *Antiviral Res* 2020;177:104762.
  - 36 Colson P, Rolain J-M, Lagier J-C, *et al.* Chloroquine and hydroxychloroquine as available weapons to fight COVID-19. *Int J Antimicrob Agents* 2020;55:105923.
  - 37 Wang M, Cao R, Zhang L, *et al.* Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* 2020;30:269–71.
  - 38 InYao X, Ye F, Zhang M, *et al.* Vitro antiviral activity and projection of optimized dosing design of hydroxychloroquine for the treatment of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* 2020;pii: ctaa237.
  - 39 Gao J, Tian Z, Yang X. Breakthrough: chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci Trends* 2020;14:72–3.
  - 40 Gautret P, Lagier J-C, Parola P, *et al.* Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *Int J Antimicrob Agents* 2020:105949.
  - 41 Available: <https://clinicaltrials.gov/ct2/results?cond=covid-19&intr=%22Antimalarials%22>
  - 42 Available: <http://www.chictr.org.cn/enIndex.aspx>
  - 43 Available: <https://www.cdc.gov/coronavirus/2019-ncov/hcp/therapeutic-options.html>

### Answers

1. A (False), B (True), C (True), D (False)
2. A (False), B (True), C (True), D (False), E (True), F (False), H (True)
3. A (False), B (True), C (False)
4. A (True), B (True), C (False), D (True), E (False)
5. A (True), B (False), C (True)